

(Affix Patient Label)
 Name.....
 Date of Birth : []/[]/[] M/ F
 NHS No : [] [] [] [] [] [] [] [] [] []
 Hospital No : [] [] [] [] [] [] [] [] [] [] [] []
 Consultant Ward.....

Post op Day 2

		Day	Night	Signature & date
Airway	<ul style="list-style-type: none"> • Laryngectomy tube care • Wean off oxygen • Humidification & Nebulisers. 			
Stoma	<ul style="list-style-type: none"> • Monitor for wound breakdown • Monitor any bleeding • Check position of Foley catheter in TEP monitoring for pressure damage & alternate side. 			
Medication	<ul style="list-style-type: none"> • Review Pain Control • Review and monitor route 			
Medical Interventions	<ul style="list-style-type: none"> • Review IV fluids if continued. • Bloods if required 			
Nutrition	<ul style="list-style-type: none"> • Review tolerance of & follow dietetic regime. 			
Swallowing	<ul style="list-style-type: none"> • Continue NBM • Encourage Saliva swallows avoiding oral suction. 			
Communication	<ul style="list-style-type: none"> • As per day 1 			
Output	<ul style="list-style-type: none"> • Following medical review remove drain if output <30mls in 24hrs. • Monitor urine & bowels 			
Physiotherapy	<ul style="list-style-type: none"> • Continue mobilization. • Monitor chest status. 			
Psychological Health	<ul style="list-style-type: none"> • Monitor mood & sleep • Refer to H&N counsellor if required. 			

(Affix Patient Label)

Name.....

Date of Birth : []/[]/[] M/ F

NHS No : [] [] [] [] [] [] [] [] [] []

Hospital No : []

Consultant Ward.....

Post op Day 10

		Day	Night	Signature & date
Airway	<ul style="list-style-type: none">As per day 9			
Stoma	<ul style="list-style-type: none">Remove stoma suturesMonitor for wound breakdown.			
Medication	If primary laryngectomy: <ul style="list-style-type: none">Monitor tolerance of oral medications.			
Medical Interventions	If salvage laryngectomy: <ul style="list-style-type: none">Contrast swallow			
Nutrition	<ul style="list-style-type: none">If oral intake safely established, stop enteral feed.Follow dietetic recommendations			
Swallow	If Salvage laryngectomy: <ul style="list-style-type: none">Review results of contrast swallow.Commence sterile water if nil surgical concerns & follow timeline as per non salvage case. If primary laryngectomy: <ul style="list-style-type: none">If tolerating level 4, progress onto level 6 food texture			
Communication	<ul style="list-style-type: none">Fit voice prosthesis providing oral intake safely establishedCommence SVR training & vocalization technique			
Output	<ul style="list-style-type: none">Consider removal of NG (if applicable)			
Physiotherapy	<ul style="list-style-type: none">As per day 8			
Psychological Health	<ul style="list-style-type: none">As per day 4			

