

Management of Sepsis in Infants and Children

Key Document code:	WAHT-TP-062	
Key Documents Owner:	Dr L Harry	Consultant Paediatrician
Approved by:	Paediatric Quality Improvement meeting	
Date of Approval:	26 th March 2021	
Date of review: This is the most current version and should be used until a revised document is in place	7 th January 2026	

Key Amendments

Date	Amendment	Approved by
February 2019	Removal of wall poster	Paediatric QIM
19 th Nov 2020	Document extended for 1 year	Paediatric QIM/Dr J West
26 th March 2021	Approved with no amendments	Paediatric QIM
9 th March 2024	Document extended for 12 months whilst under review	Dr L Harry
14 th January 2025	Document extended for 6 months whilst under review	Dr James West
7 th July 2025	Document extended whilst PEWS charts are updated and document reviewed	Dr Ezgi Seager/Dr Tom Dawson

Paediatric Intensive Care Units admit around 1000 children with severe sepsis annually. Each year severe sepsis causes more than 100 deaths in children less than four years of age. Studies suggest mortality is increased in children who receive less fluid resuscitation and inadequate vasoactive agent use. A simple guideline to encourage early aggressive management of children with severe sepsis and septic shock, along with monitoring of clinical and laboratory endpoints is aimed at significantly improving outcomes for this group of patients.

- Ensure appropriate team is involved early including paediatric consultant, anaesthetic / ICU consultant.
- Discussion with the Paediatric Intensive Care Unit (PICU) transfer team may be required at a later stage. (KIDS 0300 200 1100)
- A time line is suggested but would depend on the clinical status of the patient, area and personnel involved

1. Recognition of Sepsis

Sepsis

Defined as Systemic Inflammatory Response Syndrome (SIRS) in the presence of, or as a result of suspected or proven infection and includes the presence of two of the following four criteria, one of which must be abnormal temperature or leukocyte count:

- Core Temperature >38.5°C or <36°C
- Inappropriate tachycardia (See PEWS chart/admission documents) in the absence of external stimulus, chronic drugs, or painful stimuli; or otherwise unexplained persistent elevation over a 0.5 – to - 4 hour time period
 OR for children < 1 year old:
 bradycardia (See PEWS chart) in the absence of external vagal stimulus, β-blocker drugs or congenital heart disease; or otherwise unexplained persistent depression over a 0.5 – 1hr time period.

- Mean respiratory rate above normal for age (See PEWS chart) or mechanical ventilation for an acute process not related to underlying neuromuscular disease or the receipt of general anaesthesia.
- Leukocyte count elevated or depressed for age (not secondary to chemotherapy induced leucopenia)

Septic Shock

Defined as suspected infection with inadequate tissue perfusion exhibited by altered mental status, capillary refill time (CRT) > 2 seconds, diminished pulses, mottled peripheries (warm shock = flash CRT, bounding pulses, wide pulse pressure). This is a reflection of severe sepsis and requires aggressive resuscitation

In practice early recognition of paediatric septic shock should be defined using clinical examination not biomedical tests and diagnosis should occur before hypotension occurs as this is a pre-terminal sign. (See Severe Sepsis Screening Tool)

ANTIBIOTICS SHOULD BE GIVEN IN THE FIRST HOUR

Sepsis will be identified from the Sepsis screen on our PEWS charts on inpatients and the Sepsis screen on the admission documents ensuring all health professionals consider sepsis in the diagnosis.

2. Early Aggressive Fluid Management

Paediatric Septic shock is associated with severe hypovolaemia and children respond well to aggressive fluid management. It is important to aggressively resuscitate in the first hour as trials have demonstrated improved outcome with early fluid resuscitation with crystalloid being deemed as effective as colloid. Some studies have suggested an improved outcome for those who receive colloid (albumin) for septic shock. Rapid boluses are recommended of 20mL/kg over 5 minutes with reassessment. Initial volume resuscitation commonly requires 40-60mL/kg but can be as much as 200mL/kg. Colloid should be considered if more than 40mL/kg is required. These volumes DO NOT appear to increase the incidence of acute respiratory distress syndrome (ARDS) or cerebral oedema. It is good practice to consider intubation and ventilation once 40-60mL/kg has been administered.

3. Consider Early Inotrope Use

Inotropes should be used early. Delay in inotrope resuscitation for meningococcal sepsis led to a 23 fold increase the in adjusted odds ratio for mortality. For those children with hypodynamic shock with high systemic vascular resistance (SVR) (Cold and shut down) dopamine or adrenaline at low doses are recommended. Adrenaline has both inotropic and chronotropic activity and greater β_2 effects in peripheral vasculature allowing vasodilatation.

There is some evidence that dopamine used in hyperdynamic shock with a low SVR (warm shock) leads to a worse outcome and therefore it is advocated that low dose noradrenaline is used in this scenario.

Because of an increase in mortality with delay in time to inotrope use it is now recommended that peripheral inotropes can be used whilst central access is being obtained. The peripheral access site must be closely monitored.

4. Intubation and Ventilation

Oxygen should be optimized throughout the resuscitation. Intubation should be considered at all points in the initial resuscitation. Intubation can risk worsening of hypotension from the direct effects (myocardial depression and vasodilator effects) and indirect effects (blunting of catecholamine release)

of induction agents. Adequate fluid management and inotropic support should be ongoing before induction of anesthesia in septic shock. Ketamine is the preferred induction agent of choice for sepsis. The use of a neuromuscular blocking agent should always be considered and a short acting blocker can facilitate intubation if the provider is confident they can maintain airway patency.

5. Ongoing Monitoring and Treatment

In children at risk of absolute adrenal insufficiency or adrenal pituitary axis failure (Purpura fulminans, congenital adrenal hyperplasia, prior recent steroid exposure, hypothalamic/pituitary abnormality) and who remains in shock despite inotrope infusion, the use of intravenous hydrocortisone may need to be considered, ideally after a blood sample has been taken for subsequent determination of baseline cortisol concentration.

Further biochemical parameters requiring monitoring include

- Glucose (monitor for both hyper- and hypo-glycaemia)
- Calcium – hypocalcaemia has been shown to increase mortality
- Arterial Blood Gas (Base excess and anion gap)
- Clotting – FFP can be used to correct a prolonged INR but as infusion, not a bolus

Observations should be done every hour as a minimum but may need to be done more frequently in the initial stages of resuscitation (every 15 minutes). This will be determined by the PEWS score (see table below).

PEWS SCORE	Action & Frequency of observations
0-1	Continue monitoring – at current frequency
2	Nurse in charge must review – 2 hourly
3	Nurse in charge must review and Dr must review – 1-2 hourly
4	Nurse in charge and Dr must review Consultant informed – Hourly, continuous monitoring (minimum of saturations monitor)
5-6	Nurse in charge and Cons must review – hourly + continuous monitoring

All PEWS charts have a sepsis trigger question for PEWS to be completed at the bottom of the charts for all ages.

Please note that clinical key documents are not designed to be printed, but to be viewed on-line. This is to ensure that the correct and most up to date version is being used. If, in exceptional circumstances, you need to print a copy, please note that the information will only be valid for 24 hours

Temperature °C	39			
	38			
	37			
	36			
	35			
Sepsis 6 Triggered Y / N PTO FOR CRITERIA				
Pain Score				
Cap Refill				
Nausea Score				
Total PEWS	0-2	3-4	5-6	

Worcestershire NHS
Acute Hospitals NHS Trust

**PEWS Chart & Pain Management
0-11mths**

Attach Patient Sticker here or record

NAME: _____
NHE NO: _____
HOSP NO: _____
D.O.B: _____ Male ☐ Female ☐

Frequency of obs (every _____ hours)

Doctor/Nurse/Family contact

Respiratory Rate (Over 1 minute)

Respiratory Distress

SpO2 Saturation %

Recording O2 Saturation Method O2 Saturation

Heart Rate & Blood Pressure

BP NOT used to calculate PEWS

Consent: _____

Temperature

Sepsis 6 Triggered PTO FOR CRITERIA

Pain Score

Cap Refill

Nausea Score

Total PEWS

On the reverse of the chart is a check list to be considered each time the PEWS is calculated.

Recognition of a child at risk of Sepsis:

If a child with suspected or proven infection AND has at least 2 of the following:

- Core temperature < 36°C or > 38.5°C (38° if immunocompromised)
- Inappropriate tachycardia > 130
- Altered mental state (including: sleepiness / irritability / lethargy / floppiness)
- Reduced peripheral perfusion / prolonged capillary refill

Lower threshold of suspicion for: age < 3 months, chronic disease, recent surgery or immunocompromised

THINK: Could this child have SEVERE SEPSIS or SEPTIC SHOCK

Attach Patient Sticker here or record

NAME: _____
NHE NO: _____
HOSP NO: _____
D.O.B: _____ Male ☐ Female ☐

PAEDIATRIC EARLY WARNING SCORE - AGE 5-12 YEARS

PEWS Escalation Aid

Remember: If you feel you need more help at any time, call for help - regardless of PEWS score

0	Continue monitoring
1	Nurse in charge MUST review
2	Nurse in charge & Doctor MUST review
3	Nurse in charge & Doctor MUST review & inform Consultant
4	Nurse in charge & Consultant MUST review
5	Nurse in charge & Consultant MUST review

NURSE CONCERN

Cont Post Op Vomiting

Underlying malignant disease

1/2 hly Bronchodilators

Unstable metabolic condition ie: DKA / MCADD

Worrying / unstable condition / unrecordable BP / CPAP

NAUSEA & VOMITING SCORE

No nausea or vomiting	0
Nausea only	1
One episode of vomiting in last hour	2
More than one episode of retching/vomiting in one hour	3

VERBAL/VISUAL SELF REPORT PAIN TOOL

0	1	2	3	4	5	6	7	8	9	10
None										Severe

SCORING

CATEGORY	0	1	2
FACE	No particular expression or smile	Occasional grimace or frown, withdrawn	Frequency to constant quivering chin, clenched jaw
LEGS	Normal position or relaxed	Uneasy, restless, tense	Kicking, or legs drawn up
ACTIVITY	Lying quietly, normal position, moves easily	Squirming, shifting back and forth, tense	Arched, rigid or jerking
CRY	No cry (awake or asleep)	Moans or whimpers; occasional complaint	Crying steadily, screams or sobs, frequent complaints
CONSOLABILITY	Content, relaxed	Reassured by occasional touching, hugging or being	Difficult to console or comfort

Recognition of a child at risk of Sepsis:

If a child with suspected or proven infection AND has at least 2 of the following:

- Core temperature < 36°C or > 38.5°C (38° if immunocompromised)
- Inappropriate tachycardia > 130
- Altered mental state (including: sleepiness / irritability / lethargy / floppiness)
- Reduced peripheral perfusion / prolonged capillary refill

Lower threshold of suspicion for: age < 3 months, chronic disease, recent surgery or immunocompromised

THINK: Could this child have SEVERE SEPSIS or SEPTIC SHOCK

More Than 1/2 or Sepsis 6 triggered

Date	Time	PEWS	Sepsis 6 Y/N	Pain Score	Pain Score Agreed with Parents / Y/N	Plan	Print Name

Record Plan & Time Of Review

The Sepsis trigger must be completed at every set of observations and documented as Y/N.

If the Sepsis trigger is triggered, then how this has been escalated and what action taken must be documented on the back of the PEWS or on PEWS continuation sheet as required. Please see below for example.

Date	Time	PEWS	Sepsis	Pain score	Agreed With parents	Plan	Print name
28/09/16	2100	4	Y			Consultant to review in 15mins, rpt obs 15mins.	SarahW

Monitoring will continue on all inpatients to ensure sepsis developing on the ward will also be identified. Admission sheets also have the Sepsis 6 criteria embedded to ensure that it is considered in each

Management on the ward will include placement according to the Paediatric Side Room Priority Matrix (WAHT-PAE-129) and antibiotics according to the Paediatric Antibiotic Prescribing Guideline:
<http://www.worcsacute.nhs.uk/EasysiteWeb/getresource.axd?AssetID=81490&servicetype=Attachment>
 Intensive monitoring will include:

- Continuous pulse oximetry
- Continuous cardiac monitoring
- Temperature
- Urine output
- Assessment of conscious level
- Regular blood glucose measurements
- Base excess

Monitoring and Compliance

This section should identify how the Trusts plan to monitor compliance with and the effectiveness of this Treatment pathway. It should include auditable standards and/or key performance indicators (KPIs) and details on the methods for monitoring compliance.

WHAT?	HOW?	WHO?	WHERE?	WHEN?
<i>These are the 'key' parts of the process that we are relying on to manage risk.</i>	<i>What are we going to do to make sure the key parts of the process we have identified are being followed?</i>	<i>Who is responsible for the check?</i>	<i>Who will receive the monitoring results?</i>	<i>Set achievable frequencies.</i>
Preseptal and Orbital Cellulitis in Children Audit all cases of preseptal/orbital cellulitis that proceed to surgery	Clinical Audit/Round Table Discussion	Paediatric Clinical Governance Group		
BCG vaccine should be offered to all babies identified as being at risk as highlighted in the introduction	Audit	Obstetric Governance Committee		
Ensure babies are correctly identified and treated	Continual vigilance and occasional audit	Consultant staff with neonatal interest	Directorate audit meetings	Biannually until proven that we have high success rates

References

- 1.) *Brierley et al.* Clinical practice parameters for haemodynamic support of pediatric and neonatal septic shock: 2007 update from the American College of Critical Care Medicine.
Crit. Care. Med. 2009; 37(2): 666-688
- 2.) *Inwald et al.* Emergency management of children with severe sepsis in the United Kingdom: the results of the Paediatric Intensive Care Society sepsis audit.
Arch. Dis. Child. 2009;94:348-343
- 3.) *Goldstein et al.* International pediatric sepsis consensus conference: definitions for sepsis and organ dysfunction in pediatrics.
Paed. Crit. Care Med. 2005 Jan; 6(1):2-8
- 4.) Advanced Paediatric Life Support – The Practical Approach. 6th Ed. London: BMJ Books; 2016
- 5.) *Lampin ME et al.* Noradrenaline use for septic shock in children: doses, routes of administration and complications.
Acta Paediatrica. 2012 Sept; 101(9):e426-30

SEPSIS 6 CRITERIA WALL POSTER

Box 1: Red Flag Sepsis Criteria for Children

- *Hypotension, defined as Systolic Bloods Pressure $< 2 \text{ SD for age, (age in yrs} \times 2) + 70$
- Heart rate of > 30 above the normal upper rate limit for age (see Box 2)
- Blood gas lactate $> 4 \text{ mmol/L}$
- Prolonged capillary refill > 5 seconds
- Pale/mottled/ashen/blue or non-blanching (purpuric) rash
- Oxygen needed to maintain saturations $> 92\%$
- Respiratory rate $> 60 \text{ min}^{-1}$ or > 5 below the normal, or grunting
- AVPU = V, P or U
- Parents report excessively dry nappies, lack of response to social cues, significantly decreased activity or weak, high-pitched or continuous cry

Box 2: Age-specific RED FLAG parameters for vital signs and laboratory variables

	Heart Rate, Beats/Min	Heart Rate, Beats/Min	Respiratory Rate	Leukocyte Count	Systolic Blood Pressure
	Tachycardia	Bradycardia	Breaths/Min	Leukocytes $10^3/\text{mm}^3$	mmHg
Birth to 1 week	> 180	< 100	> 50	> 34	< 60
1 week to 1 month	> 180	< 100	> 40	> 19.5 or < 5	< 60
1 month to 1 year	> 180	< 90	> 34	> 17.5 or < 5	< 70
2 to 5 years	> 140	NA	> 22	> 15.5 or < 6	$< *$
6 to 12 years	> 130	NA	> 18	> 13.5 or < 4.5	$< *$
13 to 18 years	> 110	NA	> 14	> 11 or < 4.5	$< *$

Box 3: High Certainty of Sepsis

High Certainty of Sepsis
Respond with Paediatric Sepsis 6:

Complete all elements within 1 hour

1. Give high flow oxygen

2. Obtain IV / IO access & take blood tests

- a. Blood cultures
- b. Blood glucose – treat low blood glucose
- c. Blood gas

3. Give IV or IO antibiotics

- Broad spectrum cover as per local policy

4. Consider fluid resuscitation

- Aim to restore normal circulating volume and physiological parameters
- Titrate 20ml/kg Isotonic Fluid over 5 -10min and repeat if necessary
- Caution with fluid overload:
Examine for crepitations & hepatomegaly

5. Involve Senior Clinicians / Specialists early

6. Consider Inotropic support early

- If normal physiological parameters are not restored after $\geq 40\text{ml/kg}$ fluids
- NB Adrenaline or Dopamine may be given via peripheral IV or IO access

High Certainty of Sepsis

- Give High Flow O2
- Obtain IV/IO Access & take bloods
- Give IV or IO antibiotics
- Consider fluid resuscitation
- Involve Senior Clinicians / Specialists Early
- Consider Inotropic Support

Insert sticker in the notes to record

**For every hour that a child remains in Septic Shock
the mortality risk doubles.**

ULN = Upper limit of normal

ICE = Electronic results reporting system

PEWS = Paediatric Early Warning Score

Consider Early Inotropes:

For central access see KIDS Drug calculator.

<http://kids.bwc.nhs.uk/healthcare-professionals-2/drug-calculator/>

Peripheral administration whilst awaiting central access should be closely monitored:
Dopamine and Dobutamine peripheral strengths are available on the KIDS Calculator.

Noradrenaline

Noradrenaline 0.02mg/ml strength could be started at a dose of 0.1 micrograms/kg/min and titrated up to a maximum of 5micrograms/kg/min whilst gaining central access.

A good cannula or intraosseous needle should **only** be used for this inotrope infusion.

Adrenaline

Adrenaline 0.01 – 1.0 microgram/kg/min can be used the same strength as peripherally while obtaining central access.

A good cannula or intraosseous needle should **only** be used for this inotrope infusion.

If blood pressure drops check the cannula site before increasing the infusion rate.

SEPSIS SIX		Time Completed (24hr clock) & initials	Reason not don result
Sepsis Diagnosis Time:			
1. Oxygen: high flow 15l/min via non-rebreathe mask. Maintain SPO2 94% or more			
2. IV/IO Access Blood Cultures Blood glucose Blood Gas inc Lactate		B/C: BM: B/G	
3. IV/IO Antibiotic as per trust guidelines – Antibiotic given:			
4. IV/IO Fluids: 20mL/kg of 0.9% Saline STAT (10mL/kg alliquots if cardiac patient). Caution with fluid overload . Maintain STRICT Fluid Balance			
5. Involve senior clinicians early/Specialists Early (KIDS)			
6. Consider inotropic support early			