

Remifentanil Patient Controlled Analgesia Guideline

Owner Ramy Labib Joanna Marriott	Consultant Anaesthetist Consultant Anaesthetist
Approved by Obstetric Governance Meeting on:	21 st November 2025
Approved by Medicines Safety Committee on:	13 th January 2026
Review Date: This is the most current document and should be used until a revised version is in place.	21 st November 2028

Introduction

Neuraxial analgesia is the gold standard for pain relief in labour and well established across the UK¹. However, it may not always be possible or preferred. In these circumstances alternative forms of pharmacological analgesia can be considered, including remifentanil patient controlled analgesia (PCA).

This guidelines is for the use by the following staff groups

Anaesthetists
Midwives
Obstetricians
Midwife Support Workers
Anaesthetic theatre staff

Key Amendments

Date	Amendment	By
March 2015	New guideline	P Lo J Marriott
	Changed drug preparation from 2milligram remifentanil in 100ml bag to 1milligram remifentanil in 50ml syringe.	J Greenwood
20 June 2021	Major rewrite of guideline	M van Velze T Brunning J Greenwood
November 2025	Update to include diluting 2milligram of remifentanil in 1milligram is unavailable	Rami Labib Keith Hinton

**Obstetric Pathways
WAHT-TP-094**

CONTENTS

INTRODUCTION	3
OBJECTIVES	3
DUTIES AND RESPONSIBILITIES	3
Anaesthetist	3
Midwife	3
INDICATIONS	4
PREREQUISITES	4
CONTRAINDICATIONS	4
PROCEDURE	5
Patient preparation	5
Equipment needed	5
Drug preparation, administration and prescription	5
General points	6
Discontinuation of remifentanil PCA	6
MONITORING	7
Logistics and staff	7
Observations	7
Pain scoring:	7
Sedation scoring:	7
Indications to call the anaesthetist:	7
POINTS OF SAFETY	8
Appendix A: Remifentanil Patient Controlled Analgesia (PCA) - Patient Information Sheet	9
Appendix B: Remifentanil PCA Proforma for labour	11
Appendix C: Emergency Management	13
Appendix D: Patient information Infographic	16

INTRODUCTION

Remifentanyl is an ultra-short acting intravenous opioid. It is a hundred times more potent than morphine, with a rapid onset and peak effect within about 80 seconds. Its rapid offset is due to the deactivation by abundant non-specific plasma and tissue esterases, which also makes it non-cumulative. It freely crosses the utero-placental interface, but also undergoes rapid foetal elimination. The side effect profile is similar to other opioids, including an increased risk of respiratory depression when compared to pethidine and the use of an epidural¹. Therefore, the safety measures and other side effects will be discussed in further detail.

Despite remifentanyl proving to provide inferior analgesia compared to an epidural, it showed high maternal satisfaction when used. Modes of delivery, Apgar scores and adverse effects such as nausea, vomiting and pruritus were comparable between the two methods¹.

The RESPITE trial compared intramuscular (IM) pethidine with remifentanyl PCA and demonstrated higher patient satisfaction, lower pain scores with significantly fewer women converting to epidurals and fewer instrumental deliveries with remifentanyl². Apgar scores were similar for the two drugs³.

Even though remifentanyl PCA is seen as a safe and effective option of analgesia for labour and widely used in maternity units across the world, its use in this manner is currently unlicensed.

OBJECTIVES

- To summarise the indication, pre-requisites and contra-indications to remifentanyl PCA
- To explain the procedure of initiating a remifentanyl PCA, including drug preparation
- To ensure appropriate monitoring of the patient using a remifentanyl PCA
- To ensure safe disposal of any unused drug after use, in line with legal requirements of management of controlled drugs

DUTIES AND RESPONSIBILITIES

Anaesthetist

- Assess the suitability of the patient for a remifentanyl PCA
- Consent the patient
- Start remifentanyl PCA proforma
- Check the drug with the midwife or ODP and prepare the remifentanyl
- Set up the pre-programmed PCA pump
- Explain to the patient how to use the PCA
- Remain in the room for the first five administrations
- Prescribe the PCA, oxygen, naloxone, ondansetron and omeprazole.
- Change the remifentanyl syringes when required

Midwife

- Provide patient with remifentanyl leaflet prior to informing anaesthetist
- Check and sign out drug with anaesthetist
- Monitor the patient as per protocol
- Ensure patient receives continuous one to one care
- Alert the anaesthetist of significant side effects
- Dispose of residual drug and document in controlled drug book as per trust guideline and legal requirements.

Obstetric Pathways
WAHT-TP-094**INDICATIONS**

- **Contra-indications to epidural analgesia**
 - Patient refusal
 - Local infection or systemic sepsis
 - Coagulopathy
 - Therapeutic anticoagulants
 - Idiopathic thrombocytopenia ($<75 \times 10^9/L$)⁴
 - Secondary to acquired pathology: Pre-eclampsia (PET) with platelets $<100 \times 10^9/L$ within the last six hours⁴, Haemolysis Elevated Liver Enzymes and Low Platelets (HELLP) syndrome, cholestasis with INR > 1.4 within the last 24 hours⁴, Immune Thrombocytopenic Purpura (ITP), massive haemorrhage, sepsis, Disseminated Intravascular Coagulopathy (DIC) etc.
 - Inherited condition: haemophilia, von Willebrand's disease
 - Raised intracranial pressure
 - Technical difficulties/Structural abnormalities of the back
 - Spinal metastases
 - Previous spinal surgery (judged on individual case basis)
 - Abnormal spinal anatomy e.g. severe kyphoscoliosis
 - Some neuropathies/altered neurology in the lower limbs
- **Failed epidural analgesia**
- **Maternal choice**
 - Given prerequisites are met and no contra-indications (see below).
 - In case of low staffing levels, priority should be given to patients with contra-indications to regional anaesthesia.
 - In conditions where an epidural is advantageous for medical or obstetric reasons, including, but not limited to pre-eclampsia, extreme BMI (BMI > 45) and multiple pregnancy^{5,6} patients should have in-depth counselling on the risks and benefits of an epidural vs remifentanyl PCA and consultant discussion is advised in these cases.

PREREQUISITES

- $> 36/40$ gestation
- Singleton live pregnancy
- Intending vaginal birth
- Be in established labour with regular contractions and at least 3-4cm cervical dilatation

CONTRAINDICATIONS

- Patient refusal
- Allergy to opioids
- Other opioids administered within the preceding four hours
- Resting SpO₂ $<95\%$ or severe respiratory disease
- Intra-uterine death (offer morphine PCA or epidural if not contra-indicated)
- Evidence of foetal distress
- Intravenous drug users
- Insufficient monitoring or staffing levels
- Inability of patient to use the PCA system

**Obstetric Pathways
WAHT-TP-094**

PROCEDURE

Patient preparation

- The patient should ideally read the Remifentanil Information Leaflet (appendix A) prior to having a discussion with an anaesthetist.
- Use the remifentanil PCA proforma (appendix B)
- The patient should be consented for this therapy and consent should include
 - Explanation of the drug itself
 - Reasons why an epidural is contra-indicated, if so OR that an epidural can be requested at any time during the use of the PCA, however it will be disconnected prior to epidural insertion
 - Advantages of remifentanil over other opioids
 - Possible side effects:
 - Drowsiness
 - Dizziness
 - Nausea
 - Itching
 - Reduced respiratory rate and transient, usually self-limited lowered oxygen saturations with 2-3 in 10 patients requiring additional oxygen via nasal specs
 - Unlicensed nature of the therapy
- Explain/show how to use the PCA with important points being:
 - Timing of administrations, which should be before or at the start of a contraction to ensure the peak drug effects coincides with the peak of the contraction. Late pressing of the button will reduce effectiveness and increase the risk of side effects.
 - ONLY the patient can press the button (not the midwife or birthing partner)

Equipment needed

- Dedicated 20G/22G intravenous cannula
- Additional large bore cannula for administration of fluids/other drugs
- Pre-programmed remifentanil PCA pump – labelled ‘Maternity’
- Intravenous (IV) extension set with an anti-syphon and anti-reflux valve
- 50ml syringe
- 50ml normal saline
- 1milligram remifentanil ampule
- Nasal specs for oxygen
- Self-inflating bag valve mask/Ambu bag present in room
- Naloxone (400microgram) ampule with 2ml syringe, drawing-up needle and label present in room (does not need to be drawn up)

Drug preparation, administration and prescription

- Remifentanil is stored in the controlled drug cupboard on delivery suite.
- Dilute 1milligram Remifentanil in 50ml of 0.9% Sodium Chloride (20µg/ml) in a 50ml syringe
- Once mixed, this solution is stable for 24 hours at room temperature.
- Label the syringe appropriately.
- Attach the IV extension set and prime.
- Load the syringe into the remifentanil PCA pump and lock the box.
- There are three pre-programmed regimes, with different bolus doses of remifentanil, a two-minute lockout period and no background infusion:
 - Regime A: 1ml bolus (20microgram)
 - Regime B: 1.5ml bolus (30microgram)
 - Regime C: 2ml bolus (40microgram)
- Start with regime B (30microgram)

Obstetric Pathways
WAHT-TP-094

- Use the dedicated 20G/22G cannula and connect the PCA giving set directly to this – **do not use a Y-connector/clave. This cannula should not be used for other medications, IV fluids or IV flushes.**
- A pulse oximeter should be attached before the PCA is started and should remain in place for the duration of the PCA. Set the alarm at 95% and nasal specs oxygen should be started if saturations fall below that.
- The anaesthetist should remain in the room for the first five presses.
- If the patient is managing to synchronise her contractions and PCA doses well and pain relief is still not adequate after 15 minutes, increase to regime C (40microgram). On the other hand, if excessive sedation or respiratory depression, go to regime A (20microgram) – NEVER CHANGE THE DRUG CONCENTRATION.
- Prescribe the following:
 - On the proforma: Remifentanil PCA
 - On the drug chart: Naloxone prn
Oxygen
Ondansetron
Omeprazole/Antacid
- If 1milligram Remifentanil is not available, dilute 2milligram Remifentanil in a 100 ml bag of 0.9% Sodium Chloride (20microgram/ml). Use 50 ml as described above, and discard the rest of the bag appropriately. We currently have no safe mechanism to store the rest of the drug, and the potential risks outweigh any benefits.

General points

- Mothers can only drink clear fluids once a remifentanil PCA is started.
- Entonox and TENS (transcutaneous electrical nerve stimulation) may still be used in addition to a remifentanil PCA.
- No other opioids should be administered in conjunction to a remifentanil PCA.
- An epidural can be requested at any stage during the use of a remifentanil PCA, if not contra-indicated. However, the PCA must be disconnected prior to epidural insertion.
- Prescribe an antacid, especially if the patient has a history of reflux or when a remifentanil PCA is used because regional anaesthesia is contra-indicated and the patient is at high risk of requiring delivery by caesarean section.
- Breastfeeding can be commenced immediately after delivery AND discontinuation of the remifentanil PCA⁷.

Discontinuation of remifentanil PCA

- The PCA can be used for the duration of labour and can be continued during perineal repair.
- Once the remifentanil PCA is no longer required, it should be disconnected, and the cannula removed WITHOUT flushing it.
- The midwife is responsible for safe disposal of the remaining drug, as per trust guidelines, with the total amount administered and disposed of documented in the controlled drug book.

MONITORING

Logistics and staff

- Ensure one to one midwifery care for the duration of the PCA by a midwife trained in the care of a woman with a remifentanil PCA and Intermediate Life Support (ILS).
- An anaesthetist is to be present in the room for the administration of the first five doses.
- The anaesthetist should remain on the delivery suite whilst remifentanil is being used.

Observations

- All observations to be documented on the remifentanil PCA proforma AND on Badgernet.
- Record a set of baseline observations (HR, BP, SpO₂, RR, CTG) prior to starting a remifentanil PCA
- Saturations should be monitored continuously for the duration of the remifentanil PCA.
- HR, BP, RR, saturations, pain and sedation scores should be documented every five minutes for the first 15minutes, then half hourly until delivery, unless more frequent observation is required.
- The BP cuff should be placed on the arm which does not have the cannula attached to the remifentanil PCA.
- The integrity of the cannula, number of attempts/successful boluses and remaining volume should be documented hourly.

Pain scoring:

- 0 = Pain free
- 1 = Mild pain
- 2 = Moderate pain
- 3 = Severe pain

Sedation scoring:

- A = Alert
- V = Voice (responds to)
- P = Pain (responds to)
- U = Unresponsive

Indications to call the anaesthetist:

- Sedation scores P or U on the AVPU scale
- RR < 8 breaths/minute
- Two episodes of apnoea (>10 seconds) requiring verbal stimulus to breath
- Saturations < 90% despite 2L oxygen via nasal specs
- Maternal HR <50bpm
- Systolic BP decrease by >25% baseline or < 90mmHg
- Evidence of foetal distress as part of the normal response to such a situation and especially if any of the above features present.
- The patient does not want to continue with the remifentanil PCA
- Difficulty achieving analgesia
- There is a problem with the PCA pump

In the event that the obstetric anaesthetist is in theatre out of hours, please do not hesitate to call the fourth on (703) or first on (700) to help troubleshoot.

POINTS OF SAFETY

- Use of a dedicated 20G/22G cannula for remifentanil PCA with no Y-connector.
- PCA cannula not to be used for any other medications, fluids or flushes.
- ONLY use a pre-programmed maternity PCA pump and DO NOT CHANGE the remifentanil concentration.
- ONLY the patient can press the PCA button (not the midwife or birthing partner).
- No other opioids should be administered in conjunction to a remifentanil PCA.
- Continuous one to one midwifery care for the duration of the PCA.
- Continuous oxygen saturation monitoring while using the PCA.
- Additional oxygen administration via nasal specs, if needed.
- Mothers can only drink clear fluids while using a remifentanil PCA.
- Ensure an Ambu bag and an ampule of naloxone are present in the room.
- On discontinuation of the PCA, remove the cannula without flushing it.

Appendix A: Remifentanil Patient Controlled Analgesia (PCA) - Patient Information Sheet

What is a Remifentanil PCA?

Remifentanil is an ultra-short acting potent morphine-like substance. It is given for pain relief in labour via a dedicated cannula into your vein with a specific pre-programmed pump controlled by yourself and is therefore called 'patient controlled analgesia' or PCA. It works very quickly and also wears off quickly, meaning that it is good for the nature of labouring pains. Although it cannot provide complete pain relief, many women find it very beneficial.

Who can use a Remifentanil PCA?

Most women are able to use a remifentanil PCA. It is a useful method of pain relief when an epidural is not preferred or cannot be used, for reasons such as infection and bleeding problems, abnormalities of the spine or difficulty of insertion.

There are a few circumstances when a remifentanil PCA should not be used. These include if you have an allergy to morphine-like medicines, if you have a severe breathing problem, if pethidine or other morphine-like medicines have been used within the last four hours, and if there is insufficient staff available to monitor you whilst the remifentanil PCA is being used.

What are its side effects?

Side effects wear off quickly after stopping the PCA and may include:

- Drowsiness
- Dizziness
- Nausea
- Itching
- Slowing of breathing rate
- Decreased oxygen saturation levels

How do we ensure it is safe to use?

Safety features include

- The use of specifically programmed pump
- Continuous presence of your midwife in the room
- Continuous monitoring of your oxygen saturations
- The administration of additional oxygen via nasal specs if needed
- ONLY YOU are allowed to press the button, not the midwife or your birthing partner

What are the effects on my baby?

Even though remifentanil crosses the placenta to reach the baby, it is also rapidly deactivated in their blood and has therefore been demonstrated to be safe. It has no additional effects than those of pethidine and these effects are shorter lived.

How do I use a remifentanil PCA?

The anaesthetist will explain this to you. The pump is programmed to give you a small and exact amount of the remifentanil when you press a button. It only allows a set number of administrations in a given hour and therefore reduces the risk of having too much of it. The effect of remifentanil can be felt within 30 seconds and will wear off after a few minutes. As it does not work immediately, the button should be pressed when a contraction starts, NOT when the contraction is at its most painful. This allows time for it to reach its full effect when the contraction is at its worse and wear off as the contraction does. Pressing the button late will reduce the effectiveness and can increase the risk of side effects. We require you to only drink clear fluids once you start using a remifentanil PCA.

When can I ask for a Remifentanil PCA?

It can be used anytime during established labour. If you decide to use a remifentanil PCA, your midwife will contact an anaesthetist to discuss the matter further and to answer any questions you have. It shouldn't take too long to set up the pump and once it is connected to the cannula in your vein, it will be ready to use.

Can I use other pain relief medicines with a Remifentanil PCA?

Yes, Entonox (aka 'gas and air') and TENS machines can be still be used with a remifentanil PCA, if required. However, you won't be able to use other morphine-like medicines, such as pethidine or codeine.

If I use a Remifentanil PCA, can I request an epidural if I want one?

If an epidural is not contraindicated, then yes, you can opt for one at any time. You will need to be assessed by the anaesthetist again, as is routine for all epidurals. The remifentanil PCA must be discontinued prior to insertion of the epidural.

Is it safe to breastfeed my baby if I use a remifentanil PCA during my labour?

Yes, remifentanil is broken down by the body extremely quickly. Once you have stopped using the PCA, very little remifentanil will be in your system after 5-10 minutes. You can start breastfeeding immediately after delivery and discontinuation of the PCA.

Appendix B: Remifentanil PCA Proforma for labour – FRONT PAGE

Affix patient label here or record	
Name:	
NHS no:	
Hosp no:	
DOB:	M/F

REMIFENTANIL
PATIENT
CONTROLLED
ANALGESIA
(PCA)

Anaesthetist: _____
Date: ___/___/___ Time: ___:___

Prior analgesia:
None N2O Pethidine Other

Indication:
Regional contra-indicated: _____
Maternal request

PMH/Obstetric history:

Drug history:

Prerequisites:

<input type="checkbox"/>	>36/40 gestation
<input type="checkbox"/>	Singleton, live pregnancy
<input type="checkbox"/>	Established labour
<input type="checkbox"/>	No opioids in last 4hours
<input type="checkbox"/>	Baseline sats >95%
<input type="checkbox"/>	No opioid allergy

Consent:

<input type="checkbox"/>	Leaflet seen
<input type="checkbox"/>	Drowsiness
<input type="checkbox"/>	Dizziness
<input type="checkbox"/>	Nausea
<input type="checkbox"/>	Itching
<input type="checkbox"/>	Reduced RR
<input type="checkbox"/>	Need for O2 (1/10)
<input type="checkbox"/>	Incomplete analgesia
<input type="checkbox"/>	Unlicensed use

Equipment: _____ (20/22G)

<input type="checkbox"/>	Dedicated cannula
<input type="checkbox"/>	Additional cannula
<input type="checkbox"/>	Maternity PCA pump
<input type="checkbox"/>	Giving set
<input type="checkbox"/>	50ml syringe
<input type="checkbox"/>	50ml saline
<input type="checkbox"/>	1milligram remifentanil
<input type="checkbox"/>	Nasal specs
<input type="checkbox"/>	Self-inflating bag
<input type="checkbox"/>	Naloxone ampule

Prescription:

DATE	DRUG	Volume	PCA Lock Out Time	PCA Bolus Dose (microgram)	Signature/ Name of prescriber	Time started	Time ended
	Remifentanil 1milligram in 50ml NaCl 0.9%						
	Remifentanil 20microgram/ml	50ml	2min	Regime A 20microgram			
	Remifentanil 20microgram/ml	50ml	2min	Regime B 30microgram			
	Remifentanil 20microgram/ml	50ml	2min	Regime C 40microgram			

Date	Time	Syringe commenced/Changed
		Prepared
		Checked

- Syringe labelled
- Oxygen prescribed
- Naloxone prescribed
- Ondansetron prescribed
- Omeprazole prescribed

Appendix B2: Remifentanil PCA Proforma for labour – REVERSE PAGE

Affix patient label here or record	
Name:	
NHS no:	
Hosp no:	
DOB:	M/F

Monitoring a patient with a Remifentanil PCA
Complete PCA chart below until PCA discontinued in addition to routine observations in labour on Badgernet.

- Continuous one to one midwifery care
- Anaesthetist present for first 5 administrations

OBSERVATIONS:

Date:

	Actual time	Sats %	RR bpm	Sedation score	Pain score	PVC integrity	Attempts/ Boluses	Volume remain	Initial
Baseline									
5min									
10min									
15min									
30min									
1hour									
1h30									
2hours									
2h30									
3hours									
3h30									
4hours									
4h30									
5hours									
5h30									
6hours									

Pain score	
None	0
Mild	1
Moderate	2
Severe	3
Sedation score	
Alert	A
Voice (responds to)	V
Pain (responds to)	P
Unresponsive	U

Call the anaesthetist if:

- Sedation scores P or U on the AVPU scale
- RR < 8 breaths/minute
- Two episodes of apnoea (>10 seconds) requiring verbal stimulus to breath
- Saturations < 90% despite 2L oxygen via nasal specs
- Maternal HR <50bpm
- Systolic BP decrease by >25% baseline or < 90mmHg
- Foetal distress, especially if above features present.
- The patient does not want to continue with the remifentanil PCA
- Difficulty achieving analgesia
- There is a problem with the PCA pump

Post-delivery:

Time of delivery: _____

Obstetric Pathways
WAHT-TP-094

Mode of delivery: NVD Instrumental LSCS

Time PCA stopped: _____

Cannula removed without flush

Adverse events: Sedation Dizziness N+V Itching Decreased RR Low sats Other:

Patient satisfaction (1-10): _____

Appendix C: Emergency Management

Over-sedation

Extreme drowsiness is a significant precursor for respiratory depression.

If the patient has a sedation score of P or U on the AVPU scale i.e. not responding to voice:

- Discontinue the remifentanil PCA
- Call the anaesthetist
- High flow oxygen (15L/min) should be administered via a face mask
- Prepare naloxone 400micrograms
- Other causes of sedation should be ruled out, for example hypoglycaemia, hypoxia, hypercarbia, use of other sedating medication, cerebrovascular event, post-ictal state
- Consider changing the PCA regime to one delivering a smaller dose of remifentanil

Respiratory depression

Remifentanil is a potent respiratory depressant. Although it has a rapid clearance, extreme vigilance is required with its use.

If RR<8bpm or saturations <90%:

- Discontinue the remifentanil PCA
- Attempt to wake the patient and encourage to breath
- High flow oxygen (15L/min) should be administered via a face mask
- Call the anaesthetist
- Prepare naloxone 400micrograms
- Consider changing the PCA regime to one delivering a smaller dose of remifentanil

Respiratory arrest/Apnoea:

- If a patient is apnoeic for > 10 seconds, verbally encourage to breath and discontinue the remifentanil PCA
- If no response despite strong verbal stimulus
- Pull the emergency buzzer and/or put out a '2222' emergency call
- Administer basic life support
- Put the patient in the left lateral position
- Give 100% oxygen and assisted breaths with an Ambu bag
- Prepare naloxone 400micrograms
- Change to 15L oxygen via a non-rebreather facemask once breathing again
- Change the PCA regime to a lower dose

Bradycardia (HR<50bpm):

- Discontinue the remifentanil PCA
- Treat with glycopyrrolate or ephedrine as needed

Hypotension (SBP <90mmHg or <25% of baseline):

- Discontinue the remifentanil PCA
- Reposition the patient: head down, left lateral
- Give a 250ml fluid bolus
- If unresponsive to the above, treat with ephedrine or metaraminol/phenylephrine

**Obstetric Pathways
WAHT-TP-094**

- Look for other causes of hypotension, as haemodynamic effects with Remi PCA are often insignificant

Foetal bradycardia (FHR < 110)/Foetal distress

- Discontinue the remifentanyl PCA
- Manage as per maternity/obstetric guideline, correcting any of the above features if present.

Appendix D: Patient Information Infographic

Remifentanil Patient Controlled Analgesia (PCA)

What is it?

Remifentanil is an ultrashort acting morphine-like substance that is injected into a vein on your demand by the press of a button, which is connected to a pre-programmed pump

When is it used?

When an epidural is not safe
When an epidural cannot be sited
On maternal request

Pre-requisites?

> 36 weeks gestation
Established labour
Intention of vaginal birth
Singleton pregnancy*

When can it NOT be used?

Allergy to opioids/morphine
Any opioids received in the previous four hours
Severe lung disease
Foetal distress
Insufficient staffing

Side effects

Drowsiness
Dizziness
Nausea
Itching
Slowing of breathing
Decreased oxygen levels

Safety features

The use of a specific pump
The use of a dedicated cannula
Continuous midwifery presence
Continuous oxygen monitoring
Used of additional oxygen if needed
Only the patient can use the pump

Other common questions:

Is it safe for my baby? **YES**
Can I breastfeed as normal? **YES**
Can I still use 'gas and air'? **YES**
Can I still get pethidine? **NO**
Can I change my mind and ask for an epidural? **YES**

For more information, please don't hesitate to ask for the full patient information leaflet or to speak to an anaesthetist

REFERENCES

1. Ronel I, Weiniger CF. Non-regional analgesia for labour: remifentanil in obstetrics. *BJAed* 2019; 19(11): 357-361.
2. Wilson MJA et al. Intravenous remifentanil patient-controlled analgesia versus intramuscular pethidine for pain relief in labour (RESPITE): an open-label, multicentre randomised controlled trial. *The Lancet*. 2018, 392: 662-672.
3. Ng TKT, Cheng BCP, Chan WS, Lam KK, Chan MTV. A double-blind randomised comparison of intravenous patient-controlled remifentanil with intramuscular pethidine for labour analgesia. *Anaesthesia* 2011; 66:796-801.
4. Association of Anaesthetists of Great Britain and Ireland, Obstetric Anaesthetists' Association and Regional Anaesthesia UK. Regional anaesthesia and patients with abnormalities of coagulation. *Anaesthesia* 2013; 68: pages 966-72.
5. Jelting Y, Weibel S, Afshari A, et al. Patient-controlled analgesia with remifentanil vs. alternative parenteral methods for pain management in labour: a Cochrane systematic review. *Anaesthesia* 2017; 72: 1016–28.
6. Freeman LM et al. Patient controlled analgesia with remifentanil versus epidural analgesia in labour: randomised multicenter equivalence trial. *BMJ* 2015; 350:h846.
7. Association of Anaesthetists. Anaesthesia and sedation in breastfeeding women 2020. *Anaesthesia* 2020. doi:10.1111/anae.15179.
8. Hinova A, Fernando R. Systemic Remifentanil for Labor Analgesia. *Anesth Analg*. 2009; 109(6): 1925-9.
9. Evron S, Glezerman M, Sadan O. Remifentanil: A Novel Systemic Analgesic for Labor Pain. *Anesth Analg*. 2005; 100: 233-8.
10. Blair JM, Hill DA, Fee JP. Patient-Controlled Analgesia for Labour using Remifentanil: A Feasibility Study. *BJA*. 2001; 87: 415-20.
11. Volmanen P et al. Remifentanil in Obstetric Analgesia: A Dose-Finding Study. *Anesth Analg*. 2002; 94(4): 913-7.
12. Blair JM, Dobson GT, Hill DA. Patient Controlled Analgesia for Labour: A Comparison of Remifentanil with Pethidine. *Anaesthesia*. 2005; 60(1): 22-7
13. Thurlow JA, Laxton CH, Dick A. Remifentanil by Patient Controlled Analgesia Compared with Intramuscular Meperidine for Pain Relief in Labour. *BJA*. 2002; 88(3): 374-8.
14. Volikas I, Butwick A, Wilkinson C. Maternal and Neonatal Side-Effects of Remifentanil Patient-Controlled Analgesia in Labour. *BJA*. 2005; 95: 504.
15. Kan RE, Hughes SC and Rosen MA. Intravenous Remifentanil: Placental Transfer, Maternal and Neonatal Effects. *Anesthesiology*. 1998; 88: 1467-74.
16. Hill D. Remifentanil Patient-Controlled Analgesia should be Routinely Available for Use in Labour. *Int J Obstet Anesth*. 2008; 17: 336-9.
17. Hodgkinson P, Hughes D. Audit of Remifentanil PCA in 612 Labouring Women. *IJOA*. 2008 S10.
18. Abstracts of free papers presented at the OAA Annual Meeting 15-16th May 2008
19. Muchatuta K. Remifentanil for Labour Analgesia: Time to Draw Breath. editorial *Anaesthesia*. 2013; 68(3): 231-5.

**Obstetric Pathways
WAHT-TP-094**

Contribution List

This key document has been circulated to the following individuals for consultation:

Designation

This key document has been circulated to the chair(s) of the following committee's / groups for comments:

Committee
Dr J Greenwood (consultant Anaesthetist)