

Stroke Positioning Pathway

Long Sitting / Sitting Up in Bed

Owner:	Stephanie Flint, Clinical Lead Physiotherapist for Stroke Neuro-Rehab and Oncology
Approved by	Therapies Clinical Governance Approval Group
Approval Date	29 th March 2023
Review Date	29 th March 2026 This is the most current document and should be used until a revised version is in place

Key Amendments

Date	Amendments	Approved by:
25 th January 2023	Document extended to 30 th June 2023 whilst under review.	Dr J Trevelyan/ Benjamin Thomas
March 2023	Document re-approved for 3 years	Therapies Clinical Governance Approval Group



Ideally and where possible it is advisable not to eat and drink in this position and patients should be sat out.

HEAD: In neutral. Use pillows if necessary to support.
Try to encourage head control whilst eating and drinking and to prevent aspiration.

(Lower 2 pillows placed as a triangle, supporting both head and shoulders and upper arm. Top pillow placed across triangle to support head and upper shoulders. Head in a neutral position, try to prevent head rolling / turning to one side.)

ARM: Hemiplegic arm supported throughout its length. Lower pillow should elevate the lower arm.
Ensure the shoulders are placed forward of the body and supported with 1 or 2 pillows.
Place 1 or 2 pillows under the sound arm, to provide sufficient support.
Elbows placed in partial flexion palm facing downwards.
Wrist and fingers in neutral.

TRUNK: Should be straight and may be supported with 1 pillow to maintain lumbar curve of the back.
When using a profiling bed be aware of changes in the position of the trunk and hips.
Support with a pillow to maintain trunk position.

LEG: Both legs placed straight in bed.
Hips should be at 90 degrees.
A pillow may be placed under the knees if the hamstrings are tight.
Pillows may be used to maintain neutral position.

FOOT: Use splints if recommended to maintain good position of the foot and ankle.