

SUPRAVENTRICULAR TACHYCARDIA • 1/3

INTRODUCTION

- Supraventricular tachycardia (SVT) is the most common pathological tachycardia in newborns – can be new presentation or commenced in fetal life

RECOGNITION AND ASSESSMENT

- Sustained, accelerated non-sinus rhythm, regular and narrow-complex, originating above the level of the atrioventricular (AV) junction
- Heart rate >200 bpm
- May be 1 of 3 tachycardias:
 - atrial
 - atrioventricular nodal re-entry (AVNRT)
 - atrioventricular re-entrant (AVRT) – most common SVT in fetal and neonatal life
- Can be presenting feature of a congenital heart defect – **but** do not wait to exclude this before commencing treatment

SYMPTOMS AND SIGNS

- Can be variable with some common presentations:
 - acute onset in a baby in heart failure/shock with no previous signs and symptoms
 - fetal tachycardia during pregnancy
 - baby with irritability, poor feeding, sweating and breathlessness for hours/days before presentation
- SVT can cause reduced cardiac output due to reduced diastolic filling time
- many babies tolerate SVT well, however if tachycardia is sustained for >6 hr signs of congestive heart failure may develop, with irritability, tachypnoea and pallor

CAUSES

- No known cause in majority of babies
- Idiopathic SVT is more common in neonates than older children
- Wolf-Parkinson-White pre-excitation – only becomes visible after conversion to sinus rhythm
- Congenital heart defect, including Ebstein's and TGA

TRIGGERS

- Co-existing infections e.g. LRTI
- Manage all triggers appropriately

EXAMINATION

- Heart rate: >200 bpm
- Capillary refill
- Blood pressure
- Respiratory rate, may be normal/abnormal depending on:
 - signs of heart failure
 - co-existing respiratory conditions
 - infections
- SpO₂ may be normal, low, or of poor signal in haemodynamic compromise
- Cardiovascular and respiratory examination; may be normal aside from fast heart rate
- Examine baby for other reasons of tachycardia, including pain and environmental factors e.g. pyrexia (particularly in premature baby in incubator)

INVESTIGATIONS

- 12-lead ECG to confirm SVT diagnosis in haemodynamically stable cases
 - if baby haemodynamically unstable, or if ECG not available, defibrillator can record and print rhythm strips from 3 different leads
- Once SVT terminated, perform repeat ECG to assist with identification of pre-excitation and any other underlying rhythm abnormality
- Blood gas for:
 - acid-base balance
 - electrolytes
 - ionised calcium
- Echocardiogram to assess structural anatomy and cardiac function

SUPRAVENTRICULAR TACHYCARDIA • 2/3


MANAGEMENT

Contact cardiology team as soon as baby deemed unstable

If haemodynamic compromise:

Is there circulatory shock

- Defibrillator
- Anaesthetic support



Ongoing SVT

Consider amiodarone IV and discuss with **KIDS NTS** for conference call with **paediatric**


SVT terminated

Discuss with **paediatric cardiology** for further management


While arranging cardioversion consider:

- Try vagal manoeuvres
- Obtain IV access (**central or large peripheral vein**)
- Trial of adenosine 100–300 microgram/kg IV (up to 500 microgram/kg in presence of experienced clinician)

If no haemodynamic compromise:



Consider using adenosine 100 microgram/kg and increase to 500 microgram/kg in presence of experienced clinician



• Discuss with **paediatric cardiology** for prophylactic medication

• if SVT recurs despite prophylactic medication transfer to regional centre

• Discuss further medical management and cardioversion with **paediatric cardiology**

• If haemodynamically stable and still in SVT at 6 hr consider transfer to regional centre

SUPRAVENTRICULAR TACHYCARDIA • 3/3

ADDITIONAL INFORMATION:

Adenosine

- Give via cannula into large vein in upper limb, followed by rapid sodium chloride 0.9% flush; very short half-life of 10–30 sec – must get to the heart as quickly as possible
- Acts by slowing conduction time through the AV node
- Intraosseous access ineffective due to time taken for venous return
- Use 3-way tap with Luer-lock syringes; 1 syringe for adenosine and 1 for sodium chloride 0.9% flush

Never test cannula by aspirating blood into syringe with adenosine before injection – will lead to breakdown of adenosine

Major route of elimination via active take-up by red blood cells and vascular endothelial cells where it is metabolised

- Keep defibrillator nearby
- Capture and print rhythm, while adenosine given, via defibrillator rhythm strip or ECG recording
- Starting dose 100 microgram/kg, repeat after 2 min, if no effect increase to maximum dose of 300 microgram/kg [as in flowchart above](#)
- if experienced clinician present, maximum dose 500 microgram/kg

Vagal manoeuvres

- Cold stimulation of the trigeminal nerve (afferent branches) [leads](#) stimulation of the vagal nerve (efferent branches); slows AV node conduction
 - wrap bag of ice in towel and apply to baby's face or
 - wrap baby in towel and immerse entire head in ice-cold water for 5 sec
- Unilateral carotid massage not recommended – difficult to perform in neonates and has limited effect

DC cardioversion

- Applies direct current of electricity to the heart, synchronised to R wave of QRS complex on ECG
 - [less](#) risk of inducing ventricular fibrillation [than unsynchronised](#)
- Ideally carry out under general anaesthetic, or at least sedation
- If performed outside **NNU**, will require anaesthetic support
- Synchronised shock starting at 1 J/kg, if no response increase to 2 J/kg [as in flowchart above](#)

Chemical cardioversion:

- Discuss with [paediatric cardiology](#) if:
 - haemodynamically unstable and unresponsive to adenosine IV or DC cardioversion
 - haemodynamically stable and unresponsive to adenosine IV
- If SVT occurred in-utero consult perinatal plan and discuss with [paediatric cardiology](#)

Prophylactic medication

- When SVT has terminated, it is vital to commence medication to prevent further episodes
- Choice of prophylactic medication based on:
 - previous history of SVT (including in fetal life)
 - assessment of ECG, both in SVT and once terminated
 - cardiac function
- Discuss with [paediatric cardiology centre](#) and send ECG/echocardiogram for review

FOLLOW-UP

- Any episode of SVT – follow-up with paediatrician with expertise in cardiology/[paediatric cardiologist](#)
- Arrange:
 - baseline echocardiogram in outpatient clinic (if not already done)
 - Holter monitor