

CONJUNCTIVITIS

Conjunctivitis is a potentially blinding condition with associated systemic manifestations

RECOGNITION AND ASSESSMENT

- Conjunctival redness
- Swelling of conjunctiva and eyelids
- Purulent or mucopurulent discharge
- Vesicles on lids or adjacent skin (herpes simplex)

Differential diagnosis

- Sticky eye with blocked tear duct in which there is no inflammation of conjunctiva
- Congenital glaucoma in which there is corneal opacity
- Swelling of conjunctiva and eyelids as part of preseptal or orbital cellulitis

AETIOLOGY

Bacterial

- *Staphylococcus aureus*
- *Haemophilus influenzae*
- *Streptococcus pneumoniae*
- *Serratia spp*, *E. coli*, *Pseudomonas spp*
- *Neisseria gonorrhoeae* – typical onset aged 0–5 days: mild inflammation with sero-sanguineous discharge to thick, purulent discharge with tense oedema of eyelids
- *Chlamydia trachomatis* – typical onset aged 5–14 days: mild-to-severe swelling with purulent discharge (may be blood-stained)

Viral

- *Herpes simplex virus* (HSV)

MANAGEMENT

- 4–6 hrly eye toilet using sodium chloride 0.9%
- cooled, boiled tap water acceptable for home use

Conjunctivitis (see signs above)

- Swab all for:
 - Gram stain and bacterial culture and sensitivities
 - if other suspicions of HSV (e.g. vesicles etc.), swab for HSV PCR
 - use dry swab/moistened with viral transport media
 - place in dry tube/pot
 - use specific nucleic acid amplification test (NAAT) swab for *Chlamydia trachomatis* and *Neisseria gonorrhoeae* PCR
- Treat both eyes with:
 - frequent eye toilet as necessary
 - *Staphylococcus aureus*: fusidic acid 1% eye drops twice daily
 - otherwise: chloramphenicol 0.5% eye drops (every 2 hr for first 48 hr, followed by 3-4 times daily)
- Presentation \leq 24 hr of birth suggests gonococcal infection – inform consultant paediatrician
- If herpes suspected, begin treatment with aciclovir IV 20 mg/kg 8-hrly and aciclovir eye ointment (applied 5 times daily) while awaiting results

SUBSEQUENT MANAGEMENT

In severe non-resolving cases

- Take throat and eye swabs for viral PCR
- If herpes suspected, look for other signs of herpetic infection
- Treat suspected herpes with aciclovir IV and topical for 14 days

Conjunctivitis 2025–28

- Refer to ophthalmology
- ***Neisseria gonorrhoeae* suspected**
- Request urgent Gram stain and culture
- Assess baby for sepsis
- Swab for PCR
- Give single dose cefotaxime 100 mg/kg IV
- Refer to ophthalmology

Chlamydia

- If result positive treat:
 - oral erythromycin 14 days
- If maternal chlamydia treated successfully in pregnancy, baby does not require prophylactic treatment
- if unsure, swab baby's conjunctiva using NAAT swab
- if high risk, send bacterial swab for gonococcus and HIV antibody, HBsAg and HCV antibody from mother or baby

Gonococcal or chlamydia infection detected

- Refer mother and partner to genitourinary medicine for immediate treatment