

# INTRA-ABDOMINAL CYSTS

## INTRODUCTION

**This guideline does not apply to cystic structures which may be arising from the urinary tract**

- Antenatally detected intra-abdominal cysts include:
  - ovarian
  - intestinal duplication
  - mesenteric
  - vitello-intestinal

## SYMPTOMS AND SIGNS

- Most cysts will be asymptomatic but the following can be present:
  - abdominal pain
  - vomiting
  - abdominal distension
  - respiratory compromise
  - rectal bleeding
- Meconium pseudocyst may also be detected on an antenatal ultrasound. They will nearly always cause vomiting and abdominal distension and may be associated with underlying diagnosis of cystic fibrosis

## MANAGEMENT

### Antenatal

- Refer to/discuss appropriate place for delivery with fetal medicine unit
- Refer to paediatric surgeon for antenatal counselling

### Delivery

- In most cases, obstetric management will not alter

### Postnatal

- Resuscitate baby as normal
- Once stable, perform full postnatal physical examination (see **Examination of the newborn** guideline)

### Meconium pseudocyst

- If suspected antenatally, do not feed baby at birth
- Insert size 8 Fr nasogastric tube (NGT) immediately after birth and fix securely with tape (see **Nasogastric tube insertion** guideline)
- Empty stomach by aspirating NGT with a 10 or 20 mL syringe
  - if <20 mL aspirated, check position of tube
- Place NGT on free drainage by connecting to a bile bag
- Once stabilised, admit baby to NNU
- Replace nasogastric losses, mL-for-mL, using sodium chloride 0.9% with potassium chloride 10 mmol/500 mL IV
- Commence IV maintenance fluids (see **Intravenous fluid therapy** guideline)
- On day of birth, refer to on-call surgical team at planned place of surgery

### Other types of intra-abdominal cyst

- Unless significant abdominal distension present following birth, allow baby to feed normally and observe in postnatal ward.
- If baby well after **24 hr** with no abdominal symptoms and feeding normally then discharge
- Arrange outpatient abdominal ultrasound scan **4-6 weeks after birth**
- Consider sacrococcygeal teratoma (SCT) – could be entirely intra-abdominal and arising from pelvis, in which case discuss with paediatric surgeon

## SURGICAL REFERRAL

- Urgency will depend on clinical situation
- **Meconium pseudocyst:**
  - manage as above and refer to surgeon on day of birth
- **Symptomatic cyst:**
  - stabilise on NNU and refer to on-call surgical team on day of presentation
- **Asymptomatic cyst:**
  - abdominal ultrasound  $\leq 1$  week of birth
  - when result known, written outpatient referral to consultant paediatric surgeon
  - if arising from pelvis consider SCT and discuss with paediatric surgeon
- **Resolved cyst:**
  - ultrasound  $\leq 1$  week of birth, even if cyst appears to have resolved during pregnancy. Arrange outpatient surgical referral

## USEFUL INFORMATION

- <http://www.e-lfh.org.uk/programmes/paediatric-surgery/>