

Management of Asymptomatic Neonates at Risk of Thrombocytopenia

Key Document code:	WAHT-KD-015	
Key Documents Owner:	Dr Vivianna Weckemann	Consultant Paediatrician
Approved by:	Neonatal Guidelines Review Meeting	
Date of Approval:	11 th November 2022	
Date of Review:	11 th November 2025	
This is the most current version and should be used until a revised document is in place		

Key Amendments

Date	Amendments	Approved by
November 2022	Document approved for 3 years with no amendments	Dr Gregory/ Neonatal Guidelines Review Meeting

Introduction

This guideline is intended for use in managing well, asymptomatic newborn babies born to mothers with gestational thrombocytopenia or maternal immune thrombocytopenia (ITP).

It was developed as it was noted there is an increasing number of mothers with low platelets during pregnancy and clarification was needed on how infants born to these mothers should be managed to ensure a consistent approach.

It is important to screen these babies appropriately as a small proportion of babies born to thrombocytopenic mothers will have very low platelets, which rarely leads to significant bleeding in the neonate. These infants need to be identified, observed and treated appropriately to minimise bleeding risks.

Details of Guideline

Background

Thrombocytopenia is defined as a platelet count of $< 150 \times 10^9/L$ and is commonly seen during pregnancy; affecting around 5% of expectant mothers ¹. 70-80% of cases of maternal thrombocytopenia are due to gestational thrombocytopenia which is generally a benign condition for both mother and baby ². A diagnosis of gestational thrombocytopenia is one of exclusion and should have only been made for the mother if:

- Platelet count- greater than $70 \times 10^9/L$ but less than $150 \times 10^9/L$
- No history of thrombocytopenia outside pregnancy or 6 weeks post-partum
- No history of bleeding/bruising
- No family history of thrombocytopenia/bleeding/bruising
- Normal blood film
- No recent drugs that may cause thrombocytopenia
- No other cause suspected for the thrombocytopenia
- No evidence of pre-eclampsia ¹

Immune thrombocytopenia (ITP) is the second most common cause of isolated maternal thrombocytopenia accounting for around 3% ². Primary ITP is caused by autoimmune destruction of platelets ³. Placental transfer of maternal antibodies can result in neonatal thrombocytopenia. A rise in neonatal platelet count is usually seen by day 7 however it can take weeks or months to improve².

Please note that the key documents are not designed to be printed, but to be used on-line. This is to ensure that the correct and most up-to-date version is being used. If, in exceptional circumstances, you need to print a copy, please note that the information will only be valid for 24 hours and should be read in conjunction with the key document supporting information and/or Key Document intranet page, which will provide approval and review information.


**WAHT-KD-015
Neonatal Key Documents**

There are many other causes of maternal thrombocytopenia (Please refer to WHAT – OBS – 110 for list of alternative diagnoses) which are of varying significance to mother and baby.

Where maternal thrombocytopenia has been identified a neonatal blue form should have been completed which offers advice on the post natal management of the baby.

Please note all subsequent guidelines relate to asymptomatic neonates. If the baby has bleeding symptoms discussion with the on call consultant is needed

Management of Asymptomatic Infants Born to Mothers with Gestational Thrombocytopenia:



Please note that the key documents are not designed to be printed, but to be used on-line. This is to ensure that the correct and most up-to-date version is being used. If, in exceptional circumstances, you need to print a copy, please note that the information will only be valid for 24 hours and should be read in conjunction with the key document supporting information and/or Key Document intranet page, which will provide approval and review information.


**WAHT-KD-015
Neonatal Key Documents**

* For all babies discharged home parents should be advised to seek medical advice if baby develops bruising, bleeding, petechiae or becomes unwell

** Clinic nurses to liaise with consultant re blood results and plan for further repeats if necessary

*** If platelet count is $<50 \times 10^9/L$ discussion with the on call consultant is needed regarding CUSS and further management. Diagnosis of maternal gestational thrombocytopenia should be reviewed.

Management of Asymptomatic Infants Born to Mothers with Immune Thrombocytopenia (ITP):



Please note that the key documents are not designed to be printed, but to be used on-line. This is to ensure that the correct and most up-to-date version is being used. If, in exceptional circumstances, you need to print a copy, please note that the information will only be valid for 24 hours and should be read in conjunction with the key document supporting information and/or Key Document intranet page, which will provide approval and review information.

WAHT-KD-015
Neonatal Key Documents

* For all babies discharged home parents should be advised to seek medical advice if baby develops bruising, bleeding, petechiae or becomes unwell

** Clinic nurses to liaise with consultant re blood results and plan for further repeats if necessary

*** If platelet count is $<50 \times 10^9/L$ discussion with the on call consultant is needed regarding CUSS and further management (including possible treatment with IVIG).

References

1. WAHT-OBS-110. Worcester Hospital Guideline for the investigation of thrombocytopenia in pregnancy and the management of gestational thrombocytopenia. Cited 26.8.13
2. Gernsheimer T, James A and Stasi R. How I treat thrombocytopenia in pregnancy. Blood, 2013; 121(1):38-47
3. Neunert C, Lim W, Crowther M et al. The American Society of Haematology 2011 evidence based guideline for immune thrombocytopenia. Blood 2011; 117 (16): 4190-4207
4. Staffordshire, Shropshire and Black country newborn network guidelines. Cited at <http://www.networks.nhs.uk/nhs-networks/staffordshire-shropshire-and-black-country-newborn/documents/Thrombocytopenia%202011-13.pdf> on 15/9/13

**WAHT-KD-015
Neonatal Key Documents**

Monitoring Tool

This should include realistic goals, timeframes and measurable outcomes.

How will monitoring be carried out?

Who will monitor compliance with the guideline?

Page/ Section of Key Document	Key control:	Checks to be carried out to confirm compliance with the policy:	How often the check will be carried out:	Responsible for carrying out the check:	Results of check reported to: <i>(Responsible for also ensuring actions are developed to address any areas of non-compliance)</i>	Frequency of reporting:
	WHAT?	HOW?	WHEN?	WHO?	WHERE?	WHEN?
	Babies of mothers with thrombocytopenia have FBC (cord blood or sample from baby) on day 1	Spot checks and/or audit	as occurs	Acute paediatricians	Directorate Clinical Governance Group	two yearly
	Babies with platelets <100 have a clear plan for re-checking FBC	Spot checks and/or audit	as occurs	Acute paediatricians	Directorate Clinical Governance Group	Two yearly

Please note that the key documents are not designed to be printed, but to be used on-line. This is to ensure that the correct and most up-to-date version is being used. If, in exceptional circumstances, you need to print a copy, please note that the information will only be valid for 24 hours and should be read in conjunction with the key document supporting information and/or Key Document intranet page, which will provide approval and review information.

Contribution List

Key individuals involved in developing the document

Name	Designation
Clare Morgans	Paediatric SPR
Viviana Weckemann	Paediatric Consultant

Circulated to the following individuals for comments

Name	Designation
Dr Naeem Ahmad	Consultant Paediatrician
Dr Munir Ahmed	Consultant Paediatrician
Dr Taruna Bindal	Consultant Paediatrician
Dr Tom Dawson	Consultant Paediatrician
Dr Andrew Gallagher	Consultant Paediatrician
Dr Baylon Kamalarajan	Consultant Paediatrician
Dr Marie Hanlon	Consultant Paediatrician
Dr Kamal Nathavitharana	Consultant Paediatrician
Dr Clare Onyon	Consultant Paediatrician
Dr John Scanlon	Consultant Paediatrician
Dr Andrew Short	Consultant Paediatrician
Dr Peter van der Velde	Consultant Paediatrician

Circulated to the following CD's/Heads of dept for comments from their directorates / departments

Name	Directorate / Department
Dr Lakshmi Thirumalaikumar	Consultant Obstetrician
Dr Mark Crowther	Consultant Haematologist

Circulated to the chair of the following committee's / groups for comments

Name	Committee / group

Please note that the key documents are not designed to be printed, but to be used on-line. This is to ensure that the correct and most up-to-date version is being used. If, in exceptional circumstances, you need to print a copy, please note that the information will only be valid for 24 hours and should be read in conjunction with the key document supporting information and/or Key Document intranet page, which will provide approval and review information.