

# Enoxaparin (Inhixa®) dosing advice for VTE prophylaxis and treatment in adults

This guidance does not override the individual responsibility of health professionals to make appropriate decision according to the circumstances of the individual patient in consultation with the patient and /or carer. Health care professionals must be prepared to justify any deviation from this guidance.

#### Introduction

Enoxaparin sodium is a biological medicine that belongs to the drug class of low molecular weight heparin. Biological medications must be prescribed and dispensed according to brand. As per the formulary, Inhixa® is the brand used in Herefordshire and Worcestershire.

The purpose of this guideline is to outline the Inhixa® dosing advice for the following indications: prophylaxis dosing for medical/surgical patients, treatment dosing for DVT/PE/metallic valves and antenatal and postnatal prophylaxis dosing. This should facilitate safe and appropriate prescribing by clinicians of enoxaparin at WAHT for the listed indications.

This guideline will also provide administration guidance to nursing staff based on syringe availability, allowing for easier administration and reducing errors when performing dose calculations. The guidance encompasses dose-dependent variables such the patient's body weight and creatinine clearance.

The dosing table(s) in this document are to supersede all other relevant Inhixa® dosing guidance available.

#### This guideline is for use by the following staff groups:

This guideline is designed for use by all clinical and nursing staff managing adult patients who have been prescribed or require prescribing of prophylaxis or treatment dosing of Enoxaparin (Inhixa®) for the indications outlined in this document.

### Lead Clinician(s)

Mohima Akhtar Lead Pharmacist – Stroke &

**Thrombosis** 

Hafsah Mahmood Rotational Pharmacist

Approved by Haematology & Palliative Care on: 16<sup>th</sup> July 2025

Approved by Medicines Safety Committee on: 9<sup>th</sup> May 2025

Review Date: 16<sup>th</sup> July 2028

This is the most current document and should be

used until a revised version is in place

Enoxaparin (Inhixa®) dosing advice for VTE prophylaxis and treatment in adults		
WAHT-PHA-032	Page 1 of 12	Version 1



# Key amendments to this guideline

Date	Amendment	Approved by:
	New document approved at Medicines Safety Committee	



### ENOXAPARIN (INHIXA®) DOSING ADVICE FOR VTE PROPHYLAXIS AND TREATMENT IN ADULTS

Prophylaxis dosing for medical/surgical patients				
Body weight	Enoxaparin dose	Syringe colour		
50-99kg	≥30ml/min	40mg once a day	Yellow	
< 50kg OR CrCl 15-29ml/min		20mg once a day	Light Blue	
100kg-150kg*	≥30ml/min	40mg twice a day	Yellow	
>150kg*	≥30ml/min	60mg twice a day	Orange	

<sup>\*</sup>If CrCl 15-30ml/min use enoxaparin 20mg once a day and seek advice from consultant haematologist

If CrCl <15ml/min use unfractionated heparin calcium 5000 units SC twice a day whilst inpatient, to be discussed with consultant haematologist

if needed on discharge

Treatment doses e.g. for indications DVT*/PE/metallic valves				
Body weight (use most recent weight in pregnancy)	Twice daily dosing (1mg/kg) *	Enoxaparin syringe strength	Syringe colour	
~25 – 30kg	20mg twice a day	20mg	Light Blue	
31 – 44kg	40mg twice a day	40mg	Yellow	
45 – 54kg	50mg twice a day	60mg (Expel 0.1ml to obtain 50mg)	Orange	
55 – 64kg	60mg twice a day	60mg	Orange	
65 – 74kg	70mg twice a day	80mg (Expel 0.1ml to obtain 70mg)	Red	
75 – 84kg	80mg twice a day	80mg	Red	
85 – 94kg	90mg twice a day	100mg (Expel 0.1ml to obtain 90mg)	Black	
95 – 104kg	100mg twice a day	100mg	Black	
105 – 114kg	110mg twice a day	60mg and 60mg (Expel 0.1ml to obtain 50mg)+	Orange	
115 – 124kg	120mg twice a day	120mg	Violet	
125 – 134kg	130mg twice a day	60mg and 80mg (Expel 0.1ml to obtain 70mg)+	Orange and Red	
135 – 144kg~	140mg twice a day	120mg and 20mg	Pink and Blue	
145 – 155kg	150mg twice a day	150mg	Indigo	

If CrCl 15-29ml/min use 1mg/kg once a day

If CrCl <15ml/min, consider IV unfractionated heparin overlapped with warfarin (WAHT-HAE-019)

Extreme body weights should be discussed with your ward pharmacist and/or haematology

Antenatal and Postnatal prophylaxis dosing				
Body weight (booking in weight) Enoxaparin dose Syringe colour				
<50kg	20mg once a day	Light Blue		
50 – 90kg	40mg once a day <b>or</b> 40mg twice a day if high risk	Yellow		
91 – 130kg	60mg once a day	Orange		
131 – 170kg	80mg once a day*	Red		
>170kg	0.6mg/kg/daily*			

\*Can be given in two divided doses

#### Contra-indications to use (not an exhaustive list)

- Active bleeding
- Acquired bleeding disorder
- Inherited bleeding disorder
- Acute stroke
- Platelets <75 x 109/L
- Previous heparin induced thrombocytopenia
- Known allergy to heparin/LMWH
- Concurrent anticoagulants

#### For advice or more information please contact:

#### **Pharmacy**

- Your ward based pharmacist during working hours Monday Friday
- Weekend site team on Saturdays
- Out-of-hours for emergency advice, call the on-call pharmacist via switchboard

#### Haematology

On-call haematology via switchboard

Creatinine Clearance calculator: Creatinine Clearance (Cockcroft-Gault Equation)

Guideline for management of venous thromboembolism: WAHT-HAE-019

Thromboprophylaxis in pregnancy (VTE): WAHT-TP-093

Enoxaparin (Inhixa®) dosing advice for VTE prophylaxis and treatment in adults		
WAHT-PHA-032	Page 3 of 12	Version 1

<sup>\*</sup>Uncomplicated DVT treatment with low risk of reoccurrence can be dosed with 1.5mg/kg once a day (see next page)

<sup>+120</sup>mg and 150mg Inhixa® syringes are not graduated in 0.1ml thus not advised to have any doses expelled from them



Treatment doses e.g. for indications DVT/PE/metallic valves* in uncomplicated patients				
Body weight (use most recent weight in pregnancy)	Once daily dosing (1.5mg/kg)	Enoxaparin syringe strength	Syringe colour	
~25 – 34kg	40mg once a day	40mg	Yellow	
35 – 46kg	60mg once a day	60mg	Orange	
47 – 59kg	80mg once a day	80mg	Red	
60 – 73kg	100mg once a day	100mg	Black	
74 – 86kg	120mg once a day	120mg	Violet	
87 – 96kg	140mg once a day	100mg and 40mg	Black and Yellow	
97 – 103kg	150mg once a day	150mg	Indigo	
104 – 113kg+	160mg once a day	100mg and 60mg	Black and Orange	
114 – 126kg+	180mg once a day	100mg and 80mg	Black and Red	
>126kg+~	BMI to be calculated and 1mg/kg twice daily dosing to be used			

If CrCl 15-29ml/min use 1mg/kg once a day
If CrCl <15ml/min, consider IV unfractionated heparin overlapped with warfarin (<u>WAHT-HAE-019</u>)

<sup>\*</sup>Complicated patients (obesity, symptomatic PE, cancer, recurrent VTE or proximal thrombosis) should have 1mg/kg twice a day
†BMI to be calculated and patients who are obese should have 1mg/kg twice a day dosing

Extreme body weights should be discussed with your ward pharmacist and/or haematology

It is the responsibility of every individual to ensure this is the latest version as published on the Trust Intranet



# Monitoring

Page/ Section of Key Document	Key control:	Checks to be carried out to confirm compliance with the Policy:	How often the check will be carried out:		Results of check reported to: (Responsible for also ensuring actions are developed to address any areas of non-compliance)	Frequency of reporting:
	WHAT?	HOW?	WHEN?	WHO?	WHERE?	WHEN?
	Management of patients prescribed enoxaparin	Analysis of incident trends via directorate HATS	Annually	Directorate led HAT reviews	Directorates	Annually
	Administration of enoxaparin to by registered nurses to patients	Analysis of incidents via DATIX	Annually	Directorates	Directorates	Annually

Enoxaparin (Inhixa®) dosing advice for VTE prophylaxis and treatment in adults		
WAHT-PHA-032	Page 5 of 12	Version 1

It is the responsibility of every individual to ensure this is the latest version as published on the Trust Intranet



#### References

1. East Lancashire Hospitals NHS Trust (2024) Use of Low Molecular Weight Heparins (LMWH) Enoxaparin (Inhixa™) in Primary and Secondary Care – BEST PRACTICE GUIDELINE V8.0. East Lancashire Healthcare Economy. Available at:

https://www.lancashireandsouthcumbriaformulary.nhs.uk/docs/files/LMWH%20Best%20Practice%20Guideline%208.0%20-%20published%20%20on%2003.12.24.pdf

2. Herefordshire and Worcestershire Medicines and Prescribing Committee Low molecular weight heparins. Available at:

https://www.hereworcsformulary.nhs.uk/chaptersSubDetails.asp?FormularySectionID=2&SubSectionRef=02.08.01&SubSectionID=C100&drugmatch=3935#3935

- 3. National Institute for Health and Care Excellence (a) Parenteral anticoagulants. Available at: https://bnf.nice.org.uk/treatment-summaries/parenteral-anticoagulants/#low-molecular-weight-heparins
- 4. National Institute for Health and Care Excellence (b) Venous thromboembolism. Available at: https://bnf.nice.org.uk/treatment-summaries/venous-thromboembolism/
- 5. Newnham, D.A. and Suraweera, P. (2023) Thromboprophylaxis in Pregnancy (VTE) WAHT-TP-093.
- 6. NICE guidelines [NG158]. Venous thromboembolic diseases: diagnosis, management and thrombophilia testing. March 2020. Available from: https://www.nice.org.uk/guidance/ng158
- 7. Power, R. and Shafeek, D.S. (2023) Guideline for the Management of Venous Thromboembolism Including the management of patients receiving low molecular weight heparin WAHT-HAE-019.

#### **Contribution List**

#### **Contribution List**

This key document has been circulated to the following individuals for consultation;

Designation
Hafsah Mahmood – Rotational Pharmacist
Mohima Akhtar – Lead Pharmacist – Stroke & Thrombosis
Louise Williams – Lead Pharmacist for Women's and Paediatrics

This key document has been circulated to the chair(s) of the following committee's / groups for comments;

Committee
Pharmacy Governance Committee
Medicines Safety Committee
Haematology and Palliative Care Governance

Enoxaparin (Inhixa®) dosing advice for VTE prophylaxis and treatment in adults		
WAHT-PHA-032	Page 6 of 12	Version 1

It is the responsibility of every individual to ensure this is the latest version as published on the Trust Intranet



## **Supporting Document 1 - Equality Impact Assessment Tool**

. To be completed by the key document author and included as an appendix to key document when submitted to the appropriate committee for consideration and approval.

Please complete assessment form on next page;

It is the responsibility of every individual to ensure this is the latest version as published on the Trust Intranet







# Herefordshire & Worcestershire STP - Equality Impact Assessment (EIA) Form Please read EIA guidelines when completing this form

<u>ა</u>	Section 1 - Name of Organisation (please tick)						
	Herefordshire & Worcestershire		Herefordshire Council		Herefordshire CCG		
	STP						
	Worcestershire Acute Hospitals		Worcestershire County		Worcestershire CCGs		
	NHS Trust		Council				
	Worcestershire Health and Care	⊠.	Wye Valley NHS Trust		Other (please state)		

Name of Lead for Activity Hafsah Mahmood

Details of			
individuals	Name	Job title	e-mail contact
completing this	Hafsah Mahmood	Rotational	Hafsah.mahmood1@nhs.net
assessment		Pharmacist	
	Mohima Akhtar	Lead Pharmacist –	Mohima.Akhtar1@nhs.net
		Stroke &	
		Thrombosis	
Date assessment	26/02/2025		
completed			

### Section 2

**NHS Trust** 

Activity being assessed (e.g. policy/procedure, document, service redesign, policy, strategy etc.)	Title: Enoxaparin (Inhixa®) dosing advice for VTE prophylaxis and treatment in adults.			
What is the aim, purpose and/or intended outcomes of this Activity?	To have a readily available dosing table for Inhixa® use		sing table for Inhixa <sup>®</sup> use	
Who will be affected by the development & implementation of this activity?	×	Service User Patient Carers Visitors	× - -	Staff Communities Other
Is this:	<ul> <li>□ Review of an existing activity</li> <li>× New activity</li> <li>□ Planning to withdraw or reduce a service, activity or presence?</li> </ul>			

Enoxaparin (Inhixa®) dosing	advice for VTE prophy	laxis and treatment in adults
WAHT-PHA-032	Page 8 of 12	Version 1

Worcestershire **Acute Hospitals** 

It is the responsibility of every individual to ensure this is the latest version as published on the Trust Intranet

What information and evidence have you reviewed to help inform this assessment? (Please name sources, eg demographic information for patients / services / staff groups affected, complaints etc.	See reference list
Summary of engagement or consultation undertaken (e.g. who and how have you engaged with, or why do you believe this is not required)	See contribution list
Summary of relevant findings	

Section 3

Please consider the potential impact of this activity (during development & implementation) on each of the equality groups

The reach Foundative Group and explain your rationale. outlined below. Please tick one or more impact box below for each Equality Group and explain your rationale. Please note it is possible for the potential impact to be both positive and negative within the same equality group and this should be recorded. Remember to consider the impact on e.g. staff, public, patients, carers etc. in these equality groups.

Equality Group	Potential positive impact	Potential neutral impact	Potential negative impact	Please explain your reasons for any potential positive, neutral or negative impact identified
Age		X		
Disability		Х		
Gender Reassignment		X		
Marriage & Civil Partnerships		X		
Pregnancy & Maternity		X		
Race including Traveling Communities		X		
Religion & Belief		Х		
Sex		X		
Sexual Orientation		X		
Other Vulnerable and Disadvantaged Groups (e.g. carers; care leavers; homeless; Social/Economic		X		

Enoxaparin (Inhixa®) dosing	advice for VTE prophy	laxis and treatment in adults
WAHT-PHA-032	Page 9 of 12	Version 1

Worcestershire
Acute Hospitals

It is the responsibility of every individual to ensure this is the latest version as published on the Trust Intranet

Equality Group	Potential positive impact	Potential neutral impact	Potential negative impact	Please explain your reasons for any potential positive, neutral or negative impact identified
deprivation, travelling communities etc.)				
Health				
Inequalities (any preventable, unfair & unjust differences in health status between groups, populations or individuals that arise from the unequal distribution of social, environmental & economic conditions within societies)		X		

#### Section 4

What actions will you take to mitigate any potential negative impacts?	Risk identified	Actions required to reduce / eliminate negative impact	Who will lead on the action?	Timeframe
How will you monitor these actions?				
When will you review this EIA? (e.g in a service redesign, this EIA should be revisited regularly throughout the design & implementation)	3 years			

# <u>Section 5</u> - Please read and agree to the following Equality Statement

#### 1. Equality Statement

- 1.1. All public bodies have a statutory duty under the Equality Act 2010 to set out arrangements to assess and consult on how their policies and functions impact on the 9 protected characteristics: Age; Disability; Gender Reassignment; Marriage & Civil Partnership; Pregnancy & Maternity; Race; Religion & Belief; Sex; Sexual Orientation
- 1.2. Our Organisations will challenge discrimination, promote equality, respect human rights, and aims to design and implement services, policies and measures that meet the diverse needs of our service, and population, ensuring that none are placed at a disadvantage over others.
- 1.3. All staff are expected to deliver services and provide services and care in a manner which respects the individuality of service users, patients, carer's etc, and as such treat them and members of the workforce respectfully, paying due regard to the 9 protected characteristics.

Signature of person	Hafsah Mahmood
completing EIA	

Enoxaparin (Inhixa®) dosing	advice for VTE prophy	laxis and treatment in adults
WAHT-PHA-032	Page 10 of 12	Version 1

Worcestershi **Acute Hospitals NHS Trust** 

It is the responsibility of every individual to ensure this is the latest version as published on the Trust Intranet

r
  - 























It is the responsibility of every individual to ensure this is the latest version as published on the Trust Intranet



# **Supporting Document 2 – Financial Impact Assessment**

To be completed by the key document author and attached to key document when submitted to the appropriate committee for consideration and approval.

	Title of document:	Yes/No
1.	Does the implementation of this document require any additional Capital resources	N
2.	Does the implementation of this document require additional revenue	N
3.	Does the implementation of this document require additional manpower	N
4.	Does the implementation of this document release any manpower costs through a change in practice	N
5.	Are there additional staff training costs associated with implementing this document which cannot be delivered through current training programmes or allocated training times for staff	N
	Other comments:	

If the response to any of the above is yes, please complete a business case and which is signed by your Finance Manager and Directorate Manager for consideration by the Accountable Director before progressing to the relevant committee for approval.

Enoxaparin (Inhixa®) dosing advice for VTE prophylaxis and treatment in adults		
WAHT-PHA-032	Page 12 of 12	Version 1